

Noncommutative deformation and a topological nature of singularity Koiter

Trinh V. K.

Department of physics of polymer and crystal, MGU

E-mail : khoa@polly.phys.msu.ru

1. From the point of view of phenomenology, till now there are many models of plastic deformation. Almost in all monographs , for example [1-3,10], the following models are considered: sliding, regular, current, singular, analytical and deformation plasticity. The determinant ratio in deformation are a major problem, which was submitted for discussion. The its most simple kind was presented in the form of Genki-Nadai:

$$\begin{aligned}\sigma_{ij} &= 2G_s(T)e_{ij} & \text{at } T = \bar{T} \\ d\sigma_{ij} &= 2Ge_{ij} & \text{at } T \neq \bar{T}\end{aligned}\tag{1}$$

Where $d\sigma_{ij}$, e_{ij} - deviatory of a stress and deformation, G_s - current module of pure shift, G - the elastic module of shift, T - intensity tangent stress, $\bar{T}$ - the maximal value for a history of loading pre-eminent some value T_0 . In singularity model it looks like

$$de_{ij} = \sum_{n=1}^N df_n \frac{\partial f_n}{\partial \sigma_{ij}}\tag{2}$$

Where f_n - the current limiting surface. Here not zero are those df_n , for which on the appropriate surface f_n be performed loading, i.e.

$$\left(\frac{\partial f_n}{\partial \sigma_{ij}}\right)d\sigma_{ij} \geq 0$$

In model Iliusin it has the kind

$$\sigma_{ij} = \sum_n A_n \frac{d^n e_{ij}}{de^n} : e = \int \sqrt{de_\alpha de_\alpha}\tag{3}$$

However, if we use concept of potential of deformation as in [1-3,10-11], on definition of function H , we shall receive following determining ratio

$$\sigma_{ij} = \frac{\partial H}{\partial e_{ij}}\tag{4}$$

In our model [4], in which we consider process of plastic deformation as process of phase transition, potential of deformation (or the free energy) plays role of function Hamilton or action of system. So, if we admit Hamilton as $H = \alpha e^2 + \beta e^4 + \dots$, α, β - factors of elasticity, - deformation tensor, that as a first approximation we shall receive the formulas (1) (2) (3) (if H is expressed through the metrics of spaces). Roughly speaking, form (4) is general determining ratio for process of plastic deformation. Here there are two problems: how the fields σ, e in the time and in space is distributed ? What nature of the singularity Koiter ? We try to solve these problems consistently. For this purpose we should construct more general model of deformation.

2. As is known, all experimental data have shown, that in a plastic status the stress and deformation is strongly fluctuation. Besides the diagram $\sigma = f(e)$ for pressure - deformation is similar to the diagram $P = f(V)$ for pressure - volume for a liquid [5]. So, it is possible to consider a process of deformation as process of phase transition, or more generalized speaking, it is process of formation of structure [4]. Density of distribution of probability transition to plasticity is solution of the equation Fokker-Plank. However, if using potential character of process of deformation it is possible instead of the equation Fokker-Plank to apply the equation Schrodinger. In this approach tensor of deformation is chosen in the roles of parameter of the order. In work [1] A.A. Iliusin has presented the concept of trajectory of deformation in space E_5 . As a matter of fact the space E_5 is the one fiber in the deformation bundle [4] Further, because of fluctuation of the statuses of deformation, it is necessary for me to used " the secondary calculation " to receive a field deformations in a plastic status from the elasticity. Nevertheless it is necessary to notice, that a nature of process of plastic deformation is noncommutative nature.

So, the process of deformation $P = \{e_{ij}(x, t)\}$ is given. That is noncommutative deformation space. The deformation wave $E_{ij}(x, t)$ is noncommutative wave. For complete descriptions of distribution of this wave we have to use the space being tensor product $C(R_4) \otimes M_n$, and $C(R_4)$ - algebra of smooth function determined on usual space-time, and M_n - algebra $n \times n$ of matrixes. In this general space the matrix function $E(x, t)$ looks like (for 2-dimension case): $E(x, t) = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in C(R_4) \otimes M_2$. Now our task is construction of noncommutative deformation bundle on the $C(R_4) \otimes M_2$. This problem was performed in frameworks of noncommutative geometry Connes [6]. There are two important objects in this geometry. It is associative algebra $\mathcal{A}$ and universal algebra of the differential forms $\Omega_D^1(\mathcal{A})$ on the algebra $\mathcal{A}$. Here space of deformation $P = C(R_4) \otimes M_n$ plays role of associative algebra $\mathcal{A}$, at the same time universal algebra $\Omega_D^1(\mathcal{A})$ looks like [7]:

$$\Omega_D^1(\mathcal{A}) = \Omega_D^1(C(R_4)) \otimes \Omega_D^1(M_n)$$

It is tensor product of algebra of the differential forms above algebra of smooth function and algebra of the differential forms above algebra $n \times n$ of matrixes. In turn this algebra $\Omega_D^1(\mathcal{A})$ can be present through the direct sum of a horizontal and vertical of part

$$\Omega_D^1(\mathcal{A}) = \Omega_H^1 \oplus \Omega_V^1$$

and

$$\Omega_H^1 = \Omega_D^1(C(R_4)) \otimes M_n; \quad \Omega_V^1 = C(R_4) \otimes \Omega_D^1(M_n)$$

Thus, differential calculation on algebra $\mathcal{A}$, being in usual external differential, is chosen. More precisely, if through $Der(\mathcal{A})$ we had designated Lie algebra of derivative on $\mathcal{A}$, we shall receive

$$Der(\mathcal{A}) = (Der(C(R_4)) \otimes 1) \oplus (C(R_4) \otimes Der(M_n))$$

Differential df of an element $f \in \mathcal{A}$ also is divided for the sum

$$df = d_H f + d_V f$$

where $d_h f$ and $d_V f$ belong to Ω_H^1 and Ω_V^1 respectively. If $E_k, k \in \{1, 2, \dots, n^2 - 1\}$ is basis of algebra M_n , then $\partial_k = ad(iE_k)$ is a basis of algebra $Der(M_n)$ and $[\partial_k, \partial_l] = \Sigma C_{klm} \partial_m$. So, the basis θ^k of algebras $\Omega_D^1(M_n) \subset \Omega_D^1(\mathcal{A})$ is determined as $\theta^k(\partial_l) = \delta_l^k 1$. If through $\theta^i = (\theta^\alpha, \theta^a)$, where $\alpha = 0, 1, 2, 3; a \in \{1, 2, \dots, n^2 - 1\}$, the basis of algebra $\Omega_D^1(\mathcal{A})$ is designated, then the basis in $Der(\mathcal{A})$ will be as $e_i = (e_\alpha, e_a)$ and $e_\alpha = e_\alpha^\mu \partial_\mu$ are 4-generators for algebra $C(R_4)$, and $e_a = ad(E_a)$ is a basis of $Der(M_n)$. In this case θ^i accepts a kind

$$\theta^\alpha = \theta_\mu^\alpha dx^{-\mu}, \quad \theta^a = E_b E^a dE^b$$

Differential d_h and d_V will accept a kind

$$d_H f = e_\alpha f \theta^\alpha, \quad d_V f = e_a f \theta^a, \quad f \in \mathcal{A}$$

Besides there is an canonical element $\theta \in \Omega_D^1(M - n)$ determined as $\theta = E_k \theta^k$. Within the framework of geometry Connes, noncommutative bundle is right, or left $\mathcal{A}$ -module, or is more exact, Hermit $\mathcal{A}$ -module. On this module following connection is defined [7]

$$\omega = A + \chi$$

and A - it is an element of Ω_H^1 and χ - an element of Ω_V^1 . In turn, the element χ is divided for the sum

$$\chi = \theta + \phi$$

because of the fact, that $\Omega_D^1(M_n) \subseteq \Omega_V^1$. Here ϕ is a field Higgs. If through $\mathcal{U}$ it displays the group of gauge maps, then an element ω, A, θ, ϕ will be transformed under action $g \in \mathcal{U}$ by a rule

$$\omega' = g^{-1} \omega g + g^{-1} dg \tag{5}$$

$$A' = g^{-1} + g^{-1}d_H g, \quad \theta' = \theta, \quad \phi' = g^{1-}\phi g$$

The intensity of a field of deformation is determined through the 2-form Ω

$$\Omega = d\omega + \omega^2$$

Or

$$\Omega = F + D_H \phi + \Omega_V \quad (6)$$

And

$$\begin{aligned} F &= d_H A + [A, A] = \frac{1}{2} F_{\alpha\beta} \theta^\alpha \theta^\beta \\ \Omega_V &= \frac{1}{2} \Omega_{ab} \theta^a \wedge \theta^b, \quad \Omega_{ab} = [\phi_a, \phi_b] - C_{ab}^c \phi_c \\ D_H \phi &= d_h \phi + A\phi + \phi A = \theta^\alpha (e_\alpha + [A_\alpha, \phi]) \\ \phi &= \phi_\alpha \theta^\alpha, \quad A = A_\alpha \theta^\alpha \end{aligned}$$

Action of a field will be as

$$L = \frac{1}{2} (\Omega_{ij}, \Omega_{ij}) \quad (7)$$

Thus, in frameworks model [4], we shall receive probability of transition to plasticity

$$Z = \int \exp(-kL)$$

Where k - it is factor of elasticity.

3. We discuss the problem put above. For simplicity we shall be limited to consideration only pure deformation also we shall not take into account the interaction with a field Higgs. Actually, all above mentioned models were considered only in such similar situations. In these models was assumed about existence of limiting surfaces deformations (or loading). It is equivalent to the requirement of finiteness of action of deformation fields . Then, for search of a configuration of a field of

deformation, it is necessary to solve the equation, which is a condition extreme of action L ,

$$\delta L[A_\mu] = 0 \quad (9)$$

However, we have also other condition for action

$$- \int d^4x Tr[(F_{\mu\nu} + \pm \tilde{F}_{\mu\nu})^2]$$

where $\tilde{F}_{\mu\nu}$ is dual with $F_{\mu\nu}$. Because of that $Tr[F_{\mu\nu}, F_{\mu\nu}] = Tr[\tilde{F}_{\mu\nu}, \tilde{F}_{\mu\nu}]$, we receive

$$- \int d^4x Tr[F_{\mu\nu}, F_{\mu\nu}] \geq \mp \int d^4x Tr[\tilde{F}_{\mu\nu}, \tilde{F}_{\mu\nu}]$$

Is like as in work Belavin etc. [9], we are interested in importance of self-dual and self-antidual configuration of a field of deformation. Thus, one of a extreme condition of action shall write down as

$$\tilde{F}_{\mu\nu} = \mp F_{\mu\nu} \quad (10)$$

Belavin etc. used this condition for giving the instanton for gauge field. So, instead of the equation (9) we shall solve more simple the equation (10). For it we use method in [8] for search of multiinstanton in special case, when $n, r = 1$. Let the field of deformation $A = A_{-mu} dx^{-mu} \in \Omega_D^1(R_4) \otimes 1$, has a individual kind

$$A_\mu(x) = i B_{\mu\nu} \partial_\nu (\ln \phi(x)) \quad (11)$$

where $\phi(x)$ - it is a scalar function, and the matrix $B_{\mu\nu}$ looks like

$$B_{\mu\nu} = \frac{1}{2} \begin{pmatrix} 0 & +\sigma_3 & -\sigma_2 & -\sigma_1 \\ -\sigma_3 & 0 & +\sigma_1 & -\sigma_2 \\ +\sigma_2 & -\sigma_1 & 0 & -\sigma_3 \\ +\sigma_1 & +\sigma_2 & +\sigma_3 & 0 \end{pmatrix}$$

where $\sigma_1, \sigma_2, \sigma_3$ - it is matrixes Pauli. The gauge group is admitted as $SU(2)$. Setting A_μ in expression (6) that condition of self-dual (10) will

be as

$$\partial_\sigma \partial_\sigma (\ln \phi) + (\partial_\sigma \ln \phi)^2 = 0$$

or

$$\frac{\nabla^2 \phi}{\phi} = 0, \text{ where } \nabla^2 = \partial_\sigma \partial_\sigma \quad (12)$$

Using a method in [9] we receive instanton, being the solution (12)

$$\phi(x) = 1 + \frac{\lambda_1^2}{|x|^2} \quad (13)$$

and multiinstanton

$$\phi(x) = 1 + \sum_{i=1}^N \frac{\lambda_i^2}{|x_\mu - a_{i\mu}|^2} \quad (14)$$

where $a_{i\mu}, \lambda_i$ - it is any valid constant. Setting $\phi(x)$ from (12) and (13) in (11) we receive deformation wave as matrixes:

One-instanton

$$F_\mu(x) = -2i\lambda^2 B_{\mu\nu} \frac{y_\nu}{y^2(y^2 + \lambda^2)}$$

Multi-instanton

$$A_\mu(x) = -2iB_{\mu\nu} \left(\sum_i^N \frac{\lambda_i^2 y_{i\nu}}{y_i^4} \right) / \left(1 + \sum_j^N \frac{\lambda_j^2}{|y_j|^2} \right)$$

where $(y_i)_\mu \equiv (x - a_i)_\mu, i, j = 1, 2, \dots, N$

4. What real nature of sectors Koiter on a limiting surface?. With the account of solution of instanton, this problem can be easily clear. Really, using these instanton, we receive topological number Pontriagin

$$Q = -\frac{1}{16\pi^2} \int d^4x \text{Tr}[F_{\mu\nu} F_{\mu\nu}]$$

This number tell me, that space of deformation during plastic deformation will be is divided on Q of independent sectors. Thus, limiting surface, constructed in the same space, too is divided on Q of parts. It is sectors Koiter, described in works [1-3,10-11]. We see, that occurrence

of singularity Koiter has the topological reasons. Now we can speak, that process of plastic deformation is a process of birth and distribution of solitons and like-solitons. It is necessary to notice, that the defect, from the point of view of fluctuation, is one of kinds of solitons. The soliton wave is considered as the concentrated located energy, which is distributed in space without dissipation with constant velocity. Thus, the development of process of plastic deformation occurs with constant velocity and it needlessly increases a load. It will allow to understand, why a part of plasticity of the diagram stress - deformation has the horizontal form.

- [1] Iliusin A.A., Mechanics of continuous environment, MGU 1990.
- [2] Rabotnov Iu.N., Mechanics of a deformable body, M.:Sciens 1988.
- [3] Kliushnikov V.D., Physical and mathematical bases of durability and Plasticity, MGU 1994.
- [4] Trinh V.K.: kand. Thesis., MGU, 1993.
- [5] M.Ausloos: Solid Stat. Comm., 1986, vol 59, No 6, p. 401-404. 23, 1986, p. 2509-2512.
- [6] Connes A., Noncommutative differential geometry , Publ. I. H. E. S. 62, 257 (1986).
- [7] Daboit-Violette M., Kerner R., Madore J., J. Math. Phys., 32 (2), 1990.
- [8] Radzaraman R., Soliton and instanton in the quantum theory of a field, M.: the world 1985.

- [9] Belavin A. A., Polyakov A. M., Schwartz A. S., Tyupkin Yu. S.,
Phys. Lett., 1975, vol 59B, p 85.
- [10] Sedov L.I., Mechanics of continuous environments, .Sciens 1973.
- [11] Koiter W.T., Quart. App. Math., 1953, vol 11, no 3, p 350-354.